



REC'D 07 FEB 2005

Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

~~IB/2005/04363~~  
IB/04/4363

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03293281.6 ✓

**PRIORITY  
DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

R C van Dijk





Anmeldung Nr:  
Application no.: 03293281.6  
Demande no:

✓

Anmeldetag:  
Date of filing: 22.12.03 ✓  
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

PPG - SIPSY  
Z.l. la Croaix-Cadeau,  
B.P. 79  
49242 Avrillé cedex  
FRANCE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:  
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.  
If no title is shown please refer to the description.  
Si aucun titre n'est indiqué se référer à la description.)

New process for the synthesis of eneamide derivatives

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)  
revendiquée(s)  
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/  
Classification internationale des brevets:

C07D/

An Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of  
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL  
PT RO SE SI SK TR LI



## NEW PROCESS FOR THE SYNTHESIS OF ENEAMIDE DERIVATIVES.

The present invention relates to a new process for the large-scale preparation of ene-amide derivatives useful as valuable substrates for asymmetric hydrogenation reaction and hence for the synthesis of enantiomerically pure amines derivatives known as key intermediates for active pharmaceuticals.

Several methods have been described in the prior art, for example in WO 99/18065 to prepare ene-amide precursors, but these methods are clearly not very general and unsuitable for large-scale production.

The articles JOC, 1998, 63, p 6084 of the authors M. Burk and Coll. and JOC, 1999, 64(6), p 1775 of the authors X. Zhang and Coll. describe a process for ene-amide compounds synthesis comprising the reduction of oxime derivatives with iron metal in presence of acetic anhydride/acetic acid or acetic anhydride only.

The US4194050 patent describes a process for ene-amide compounds synthesis comprising the reduction of oxime derivatives with ruthenium catalyst in presence of carboxylic anhydride.

However, these processes show some limitations such as product decomposition under these conditions, use of co-solvent to facilitate product isolation, impure ene-amides which required arduous purifications and low to moderate yields.


Prior art processes are unsuitable for large-scale production of ene-amide derivatives and hence not applicable to the commercial preparation of chiral amines via asymmetric hydrogenation.

The process according to the invention presents the advantages of obtaining ene-amides in good yields, great facility of product isolation, an excellent chemical purity of product and reproducible process.

The process according to the present invention is clearly suitable for the large-scale industrial production of amine derivatives, via an asymmetric or not hydrogenation reaction. These amine derivatives, asymmetric or not, are used as intermediates for active pharmaceuticals preparation.

The present invention relates to a new process for the preparation of compounds of formula (I), comprising a hydrogenation-isomerization reaction of compound of formula (II) with an acyl derivative of formula (III) in presence of a heterogeneous catalyst as shown in scheme (I).

scheme (I) :



wherein

R<sup>1</sup> and R<sup>2</sup> and R<sup>3</sup> are independently a hydrogen atom, an alkyl, a cycloalkyl, a cycloalkylalkyl, an alkylaryl, an aryl, a heterocycle, a cyano, an acyl, an alkoxy, an aryloxy,

a carboxyl, a carbamoyl, -CONR<sup>5</sup>R<sup>6</sup> (in which R<sup>5</sup> and R<sup>6</sup> are independently an alkyl, arylalkyl or aryl group) or

-COOR5 group (in which R5 is an alkyl, alkylaryl or aryl group),

said alkyl, cycloalkyl, cycloalkylalkyl, alkylaryl and aryl groups being substituted or not with a functional group or with R5;

or R1 and R2 taken together, may form a ring (which terms includes mono-, di- and higher polycyclic ring systems);

R3 has the same meanings as R1 or R3 is a hydroxyl, imidazolyl, N-acylimidazolyl, indolyl, N-acylindolyl group or one of the groups, -COR5, -OR5, -OCOR5, -NR5R6, -NHCOR5, -N(COR5)<sub>2</sub>, wherein R5 and R6 are independently an alkyl, alkylaryl or aryl group;

R4 is a hydrogen atom, an alkyl, an aryl, an alkylaryl, said groups are substituted or not with a halogen atom as Cl, Br, or F;

Y is a hydrogen atom or R4CO group;

X is an oxygen atom or a leaving group and

m is an integer 1 or 2;

when m is 1 then X is a leaving group; when m is 2 then X is a oxygen atom.

As used herein, unless the context otherwise requires:

The term "alkyl" preferably means a straight or branched alkyl group having 1 to 20 carbons atoms such as, but not limited to, methyl, ethyl, n-propyl, isopropyl, n-butyl, iso-butyl, sec-butyl, tert-butyl optionally substituted with a functional group or with R5.

30

The term "cycloalkyl" preferably means a cycloalkyl group having 3 to 20 carbon atoms, such as, but not limited to, cyclopropyl, cyclopentyl, cyclohexyl optionally substituted with a functional group or with R5.

The term "cycloalkylalkyl" preferably means a cycloalkylalkyl group having 3-20 carbon atoms such as but not limited to cyclopropylmethyl, cyclohexylmethyl optionally substituted with a functional group or with R5.

5       The term "aryl" preferably means an aryl group having 6 to 20 carbon atoms such as but not limited to phenyl, tolyl, xylyl, cumenyl, naphthyl optionally substituted with a functional group or with an alkyl or with a fused aryl, or "aryl" means a heteroaryl group having 6 to 20 carbon atoms  
10 such as, but not limited to, furyl, thienyl, pyrrolyl, imidazolyl, pyridyl, pyrazyl, pyrimidinyl, indolyl, carbazolyl, isoxazolyl, isothiazolyl optionally substituted with a functional group or with R5 or with an alkyl or with a fused aryl.

15       The term "alkylaryl" preferably means an alkylaryl group having 6 to 20 carbon atoms such as, but not limited to, benzyl, phenethyl, naphthylmethyl optionally substituted with a functional group or with R5.

20       The term "heterocycle" preferably means a heterocycle group having 6 to 20 carbon atoms comprising one more heteroatom as O, N or S such as but not limited to pyrrolidinyl, piperazinyl, piperidyl, imidazolidinyl, piperidyl, indolinyl, said heterocycle being saturated or not, said heterocycle being optionally substituted with a  
25 functional group or with R5 or a fused aryl group.

30       The term "acyl" means preferably -COR5 (wherein R5 is as defined above or such as but not limited to acetyl, propionyl, pivaloyl, benzoyl, phenylacetyl). carboxyl, -CONR5R6 in which R5 and R6 are independently an alkyl, alkylaryl or aryl group) or -COOR5 group in which R5 is an alkyl, an alkylaryl or an aryl group,

The term "functional group" means halogen atom, or a group comprising -OH, -OR5, -CN, -COOR5, -COR5, -CONR5R6, -




OCOR<sub>5</sub>, -NH<sub>2</sub>, -NHR<sub>5</sub>, -NR<sub>5</sub>R<sub>6</sub>, -NHCOR<sub>5</sub> and -N(COR<sub>5</sub>)<sub>2</sub>, -NO<sub>2</sub>, imidazolyl, N-acylimidazolyl, indolyl, N-acylindolyl, -SH, SR<sub>5</sub>, wherein R<sub>5</sub> and R<sub>6</sub> are independently an alkyl, an alkylaryl or an aryl group,

5

The term "leaving group" means preferably one of the groups -COR<sub>5</sub>, -SO<sub>2</sub>R<sub>5</sub>, -COH, -COCCl<sub>3</sub>, -SO<sub>2</sub>F, -SO<sub>2</sub>CF<sub>3</sub>, -SO<sub>2</sub>CH<sub>2</sub>CF<sub>3</sub>, wherein R<sub>5</sub> is an alkyl, an alkylaryl or an aryl group

10

The term "ring" preferably means the formation of ring having 4 to 30 carbon atoms, such as but not limited, compounds of formula hereunder



15


wherein -R<sub>1</sub>-R<sub>2</sub>- is a methylene, dimethylene, trimethylene, tetramethylene, pentamethylene or hexamethylene linkage optionally substituted with a functional group or a fused aryl.

20

The most preferable compounds are represented by the following formula:


25

formula (IIA)



wherein  $n_1$  is an integer from 0 to 4,  $m_1$  and  $m_2$  are  
 5 each an integer from 0 to 4,  $R_7$  and  $R_8$  different or same,  
 are an hydrogen atom, a functional group, an alkyl, an aryl.





formula (IIB)



10

wherein each  $n_1$  and  $n_2$  is an integer from 0 to 4,  $Q$  is  
 an aryl, heteroaryl, cycloalkyl, heterocycloalkyl said group  
 15 are substituted or not with at least one functional group  
 preferably alpha- or beta-tetralone-oxime derivatives,  
 alpha- or beta-indanone-oxime derivatives, substituted or  
 not with a functional group.


20



Wherein R3, R7, R8 are as defined above, R9, R10 are  
 5 independently an hydrogen atom, a functional group, an  
 alkyl, an aryl.

Formula (IIC)


10



wherein n1, n2, R3 and Q are as defined above, R11 is  
 a hydrogen atom, a functional group, an alkyl, an aryl.

15

Formula (IID)



wherein n1, n2, R3, R7, R8, R9 and Q are as defined above.

Heterogeneous catalysts are based on metal like Pd, Ir, Pt, Rh, Ni catalysts preferably Ir.

The heterogeneous catalysts is used in the form of an oxide or metallic and may be supported on a suitable carrier (for example Ir/carbon or Ir/alumina).

The method how to carry out the present invention will be explained hereinafter.

The compound of formula (II) may be used as a syn-form, anti-form or a mixed-form of both.

The compound of formula (III) should be used in an amount of at least 2 molar equivalents for one molar equivalent of the oxime and may be used in a large amount as a reacting agent combined with a solvent.

The amount of the catalyst used is in the range of 0.001 to 30% mol, for 1 mol of the oxime derivative.

5 The process of the present invention is carried out in a suitable solvent. Suitable solvents are aprotic non-basic solvents such as ethers (such as but not limited to tetrahydrofuran, tetrahydropyran, diethyl ether, etc.) or aromatic hydrocarbons (such as but not limited to benzene, toluene, etc.) or carboxylic anhydrides or halogenated  
10 hydrocarbons or lower carboxylic acids or mixtures thereof.

The process of the present invention is carried out under a temperature range of -20 to 150 °C, preferably between 20 °C to 120 °C.

15

The hydrogenation of the present invention is carried out under a hydrogen pressure between 0.5 to 20 bars.

The process of the present invention is carried out  
20 for a period of time in the range of 0.5 to 24 hours.

The invention will be better understood from the experimental details, which follow.

#### Examples:

25 The present invention will be illustrated by the following examples, which will not limit the scope of the invention in any way.

#### Example 1.

30 5.5 g (0.0341 mol) of 3,4-dihydro-1H-naphtalene-2-one oxime was dissolved in 42 ml of THF. Then 9.66 ml of acetic anhydride was added dropwise. The reaction mixture is stirred at a temperature between 20-30 °C during 2 hours.

To this reaction mixture is added 0.44 g of the 5% Ir-carbon catalyts.

Then the hydrogenation is carried out at a hydrogen pressure of 6 bars and at 75 °C during 3 hours.

5        After the catalyst was filtered off, the filtrate was concentrated to dryness under reduced pressure. The residue was dissolved in 120 ml of toluene and concentrated to dryness under reduced pressure.

10       This new residue was recrystallized in a mixture of 10 ml of MTBE and 9 ml of hexane to obtain 3.82 g of the product, the compound N-(3,4-dihydro-naphtalene-2-yl)acetamide

Crude yield: quantitative / Isolated yield: 59.9- %

Chemical purity (GC): 98.95 %.

15


Example 2.

20       The reaction is carried out in the same manner as in example 1, except that 1-indanone-oxime, methoxy-6- is used as started material. The yield is 83.8 % .The chemical purity is 98.4 %.

## CLAIMS

1. A process for the production of ene-amide derivatives represented by the formula (I)

5



(I)

wherein

R1 and R2 and R3 are independently a hydrogen atom, an alkyl, a cycloalkyl, a cycloalkylalkyl, an alkylaryl, an aryl, a heterocycle, a cyano, an acyl, an alkoxy, an aryloxy, a carboxyl, a carbamoyl, -CONR5R6 (in which R5 and R6 are independently an alkyl, arylalkyl or aryl group) or -COOR5 group (in which R5 is an alkyl, alkylaryl or aryl group),

15 said alkyl, cycloalkyl, cycloalkylalkyl, alkylaryl and aryl groups being substituted or not with a functional group or with R5;

or R1 and R2 taken together, may form a ring (which terms includes mono-, di- and higher polycyclic ring systems);

20 R3 has the same meanings as R1 or R3 is a hydroxyl, imidazolyl, N-acylimidazolyl, indolyl, N-acylindolyl group or one of the groups, -COR5, -OR5, -OCOR5, -NR5R6, -NHCOR5, -N(COR5)<sub>2</sub>, wherein R5 and R6 are independently an alkyl, 25 alkylaryl or aryl group;

R4 is a hydrogen atom, an alkyl, an aryl, an alkylaryl, said groups are substituted or not with a halogen atom as Cl, Br, or F;

Y is a hydrogen atom or R4CO group;

5 X is an oxygen atom or a leaving group and

m is an integer 1 or 2;

when m is 1 then X is a leaving group; when m is 2 then X is a oxygen atom,

10 which comprise :

a hydrogenation/isomerization reaction in presence of a heterogeneous catalyst, of an oxime derivatives of formula (II)



(II)

15

wherein R1, R2 and R3 are as defined above with an acyl derivative of formula (III)



20

wherein R4, m and X are as defined above.

2. A process according to claims 1 to 2, wherein the derivative of formula (III) is used in the amount of at least 2 times per mole based on the oxime and may be used in a large amount as a reacting agent combined with a solvent.

25



3. A process according to claims 1 to 2, wherein the heterogeneous catalyst is based on metal like Pd, Ir, Pt, Rh, Ni catalyst.


5 4. A process according to claims 1 and 3, wherein the heterogeneous catalyst is used in the form of an oxide or metallic and may be supported on a suitable carrier and is used in the amount of 0.001 to 30% mole, based on the oxime derivative.

10 5. A process according to claims 1, which is carried out in a suitable solvent.

15 6. A process according to claims 1, which is carried out under a hydrogen pressure between 0.5 to 20 bars °C.

7. A process according to claims 1, which is carried out under a temperature range of -20 to 150 °C, preferably between 20 °C to 120 °C.

20 8. Ene-amide derivative of formula (I)



(I)

25 wherein

R1 and R2 and R3 are independently a hydrogen atom, an alkyl, a cycloalkyl, a cycloalkylalkyl, an alkylaryl, an aryl, a heterocycle, a cyano, an acyl, an alkoxy, an aryloxy,

5 a carboxyl, a carbamoyl, -CONR<sub>5</sub>R<sub>6</sub> (in which R<sub>5</sub> and R<sub>6</sub> are independently an alkyl, arylalkyl or aryl group) or -COOR<sub>5</sub> group (in which R<sub>5</sub> is an alkyl, alkylaryl or aryl group),

10 said alkyl, cycloalkyl, cycloalkylalkyl, alkylaryl and aryl groups being substituted or not with a functional group or with R<sub>5</sub>;

or R1 and R2 taken together, may form a ring (which terms includes mono-, di- and higher polycyclic ring systems);

15 R3 has the same meanings as R1 or R3 is a hydroxyl, imidazolyl, N-acylimidazolyl, indolyl, N-acylindolyl group or one of the groups, -COR<sub>5</sub>, -OR<sub>5</sub>, -OCOR<sub>5</sub>, -NR<sub>5</sub>R<sub>6</sub>, -NHCOR<sub>5</sub>, -N(COR<sub>5</sub>)<sub>2</sub>, wherein R<sub>5</sub> and R<sub>6</sub> are independently an alkyl, alkylaryl or aryl group;

20 R4 is a hydrogen atom, an alkyl, an aryl, an alkylaryl, said groups are substituted or not with a halogen atom as Cl, Br, or F;

Y is a hydrogen atom or R<sub>4</sub>CO group;

X is an oxygen atom or a leaving group and


25 m is an integer 1 or 2;

when m is 1 then X is a leaving group; when m is 2 then X is a oxygen atom,

30 9. Use of compounds of formula (I) as defined in claim 8 in an hydrogenation reaction, asymmetric or not, for the obtention of an amine or amide derivative of pharmaceutical interest.


## ABREGÉ

A process for the production of ene-amide derivatives represented by the formula (I)



(I)

- 5 wherein R1 and R2 and R3 are independently a hydrogen atom, an alkyl, a cycloalkyl, a cycloalkylalkyl, an alkylaryl, an aryl, a heterocycle, a cyano, an acyl, an alkoxy, an aryloxy, a carboxyl, a carbamoyl, -CONR5R6 (in which R5 and R6 are independently an alkyl, arylalkyl or aryl group) or -COOR5 group (in which R5 is an alkyl, alkylaryl or aryl group), said alkyl, cycloalkyl, cycloalkylalkyl, 10 alkylaryl and aryl groups being substituted or not with a functional group or with R5; or R1 and R2 taken together, may form a ring (which terms includes mono-, di- and higher polycyclic ring systems); R3 has the same meanings as R1 or R3 is a hydroxyl, imidazolyl, N-acylimidazolyl, indolyl, N-acylindolyl group or one of the groups, -COR5, -OR5, -OCOR5, -NR5R6, -NHCOR5, -N(COR5)2, wherein R5 and R6 15 are independently an alkyl, alkylaryl or aryl group; R4 is a hydrogen atom, an alkyl, an aryl, an alkylaryl, said groups are substituted or not with a halogen atom as Cl, Br, or F; Y is a hydrogen atom or R4CO group; X is an oxygen atom or a leaving group and m is an integer 1 or 2; when m is 1 then X is a leaving group; when m is 2 then X is a oxygen atom, which comprise : a hydrogenation/isomerization reaction 20 in presence of a heterogeneous catalyst, of an oxime derivatives of formula (II)



(II)

25

wherein R1, R2 and R3 are as defined above with an acyl derivative of formula (III) (R4CO)<sub>m</sub>X wherein R4, m and X are as defined above.

**PCT/IB2004/004363**

